microPublication BIOLOGY

11/10/2026 - *Open Access*

Complete Genome Sequences of *Mycobacterium smegmatis* Phages SVoro and PrinceCharming

Kristi M Westover[1*], Kailyn C Cabana[1*], Simon SE Cherry[1*], Norah B Kay[1*], Saachi Mehra[1*], Olivia L Morgan[1*], Ash H Olivet[1*], Samuel W Reese[1*], Sasha J Vorontsov[1*], Ella A Williams[1*], Mohammed RS Zyara[1*], Victoria J Frost[1§*]

[1]Department of Biology, Winthrop University, Rock Hill, SC, United States

[§]To whom correspondence should be addressed: frostv@winthrop.edu

[*]These authors contributed equally.

Abstract

SVoro and PrinceCharming were isolated from moist soil on the campus of Winthrop University and isolated on the host bacterium *Mycobacterium smegmatis* mc^2 155. SVoro has a myovirus morphology and a genome of 155,109 bp, while PrinceCharming exhibits siphovirus morphology and a genome of 67,429 bp. Lysogens could not be raised for SVoro or PrinceCharming, suggesting these are virulent phages. SVoro is classified in the C1 subcluster, and PrinceCharming belongs to the B2 subcluster.




Figure 1. Particle and plaque morphologies for SVoro and PrinceCharming:

Transmission electron micrographs of mycobacterium phages SVoro (A) and PrinceCharming (B). Phage lysates were negatively stained with 1% uranyl acetate, and images taken with a JEOL JEM-1230 TEM at 100kV acceleration voltage. SVoro (C) forms clear plaques (5-8 mm, n=5). PrinceCharming (D) forms relatively small, clear plaques (1-2 mm, n=4).

Description

There is a growing need for alternative treatments of antibiotic-resistant bacterial infections as they become more difficult to manage (Chinemerem Nwobodo et al., 2022). Using bacteriophages therapeutically is a promising option. Phage therapy has demonstrated positive clinical outcomes in treating bacterial infections caused by multidrug-resistant pathogens in immunocompromised patients, including after organ transplantation (Green et al., 2023), in burn wound cases (Teney et al., 2024), and in carbapenem-resistant *Pseudomonas aeruginosa* lung infections (Yang et al., 2025). In the report described here, two mycobacteriophages, SVoro and PrinceCharming, were extracted from soil near tree roots in the Hardin Garden at Winthrop University, Rock Hill, South Carolina (see Table 1 for GPS coordinates). A buffer (10% glycerol, 1 mM $CaCl_2$) was added to the samples, then shaken for 24 hours at 24° C to loosen soil particles. After settling, the liquid was filtered (0.22 µm) and enriched with *Mycobacterium smegmatis* mc^2 155 (*M. smegmatis*) in 7H9 liquid medium for four days, with shaking (250 rpm at 37° C). Following re-filtration, standard plaque assay procedures (Zorawik et al., 2024) were used to purify and amplify both phages. This involved mixing phage lysates with molten 7H9 agar and the bacterial host *M. smegmatis*, overlaying the mixture on 7H9 agar plates, and incubating for 2 days at 37° C. The plaques for SVoro were clear and relatively large (5-8 mm, n=5), while PrinceCharming's plaques were small (1-2 mm, n=4) but also clear, indicating a lytic lifecycle for both phages. This was supported by the inability of either phage to produce lysogens (Murphy et al., 2025ab). Transmission electron microscopy revealed that SVoro has a short, contractile tail (79-85 nm, n=6) typical of a myovirus morphology, with a relatively large capsid (79-87 nm diameter, n=4), whereas PrinceCharming has a long, flexible tail (221-261 nm, n=5) characteristic of a siphovirus morphology, and a smaller capsid (66-73 nm diameter, n=5).

High-titer lysates from both phages were used to extract genomic DNA with the Wizard DNA Cleanup kit (Promega), and the resulting DNA was sequenced on an Illumina NextSeq 1000 (XLEAP-P1 kit) platform, with libraries prepared using the NEB Ultra II FS kit. Raw 100 base reads were trimmed with cutadapt v4.7 (using the option: –nextseq-trim 30) (Martin 2011), and filtered with skewer v0.2.2 (using the options: -q 20 -Q 30 -n -l 50) (Jiang et al., 2014), then assembled with Unicycler v0.5.1 and Consed v29 (Wick et al., 2017; Gordon and Green 2013). Sequencing results and features of the phage genome are detailed in Table 1, which presents the predicted gene count and the phage cluster assignment based on gene content similarity (GCS) of at least 35% to phages in the Actinobacteriophage database (https://phagesdb.org/) (Russell and Hatfull 2017; Pope et al., 2017).

Bioinformatic analyses were conducted using comparative and predictive software with default settings, unless otherwise specified. Sequences were annotated with PECAAN (v20240320) (Rinehart et al., 2016), integrated with Glimmer v3.02 (Delcher et al., 2007), GeneMark v4.28 (Besemer and Borodovsky, 2005), Starterator v558 (http://phages.wustl.edu/starterator/), and Phamerator (Cresawn et al., 2011) using the Actino_draft database (v578). Additional tools included BLAST (Altschul et al., 1990) against the Actinobacteriophage and NCBI non-redundant database, and HHpred v3 (Söding et al., 2005) using PDBmmCIF70, Pfam v.37, and NCBI Conserved Domains databases. Predicted transmembrane domains were analyzed with TOPCONS v2 (Bernsel et al., 2009), DeepTMHMM v1.0.44 (integrated into PECAAN) (Hallgren et al., 2022), and SOSUI v1.11 (Hirokawa et al., 1998). Transfer RNAs were detected using Aragorn v1.2.38 (Laslett and Canback 2004), and v1.1 embedded into DNA Master (Pope and Jacobs-Sera 2018), and tRNAscan-SE v2.0.6 (Lowe and Eddy 1997).

Annotation of SVoro and PrinceCharming revealed that both genomes contain putative genes that encode proteins involved in the structure and assembly of the phage (e.g., terminase, capsid proteins, tape measure proteins, portal proteins), lysis (e.g., lysin A and lysin B), and those involved in recombination and replication (e.g., DNA resolvase, DNA primase/polymerase). Neither phage genome contains genes predicted to encode integrase or repressor functions. Exclusive use of a lytic lifestyle was supported by SVoro's and PrinceCharming's inability to create a lysogen (Murphy et al., 2025ab). SVoro contains many tRNAs in its genome (32), one tmRNA, and genes encoding putative functions such as a PnuC-like nicotinamide riboside transporter and a ThyX-like thymidylate synthase, all typical of C1 phages annotated so far (Cobb et al, 2025). PrinceCharming has the characteristic genome arrangement seen in the B2 cluster, with most genes at the right end of the genome encoded in the reverse direction, and three putative genes involved in queuosine biosynthesis conserved among subcluster B2 phages (Clermont et al., 2024).

Data availability

The complete genome sequences of phages SVoro and PrinceCharming are available in GenBank (accession numbers PV940981 and PV915860, respectively). The raw sequencing reads are available in the NCBI SRA under accession numbers SRX32748213 and SRX32748211, respectively. The Actinobacteriophage sequencing BioProject accession number is PRJNA488469.

TABLE 1 Phage sample locations and genome assembly results

Phage Name	SVoro	PrinceCharming
Sample Location (GPS)	34.9375 N, 81.030556 W	34.93714 N, 81.03081 W
Sequencing Reads	2.3M	1.9M
Sequencing coverage (X)	1443	2662
Genome size (bp)	155,109	67,429
Genome ends	Circularly permuted	Circularly permuted
GC content (%)	64.8	68.9
No. of genes	228	92
No. of tRNAs	32	0
Cluster	C1	B2

Acknowledgements: We thank the Howard Hughes Medical Institute SEA program for their continued support (especially Dan Russell and Rebecca Garlena at the University of Pittsburgh). We also thank Ms. Sigmon at the University of South Carolina (Columbia) for providing transmission electron microscopy (TEM) micrographs.

Both phages described here were collected and annotated by undergraduate students in the 2024-25 SEA-PHAGES program at Winthrop University, Rock Hill, SC.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. Journal of Molecular Biology 215: 403-410. DOI: 10.1016/S0022-2836(05)80360-2

Bernsel A, Viklund H, Hennerdal A, Elofsson A. 2009. TOPCONS: consensus prediction of membrane protein topology. Nucleic Acids Research 37: W465-W468. DOI: 10.1093/nar/gkp363

Besemer J, Borodovsky M. 2005. GeneMark: web software for gene finding in prokaryotes, eukaryotes and viruses. Nucleic Acids Research 33: W451-W454. DOI: 10.1093/nar/gki487

Clermont K, Ayuk M, Whitefleet-Smith JL, McClory S, Kirkpatrick B, Reardon S. 2024. Subcluster B1 Annotation Report. HHMI Science Education Alliance (SEA) Faculty Group, QUBES Educational Resources. http://dx.doi.org/10.25334/AGPE-VW91

Cobb ID, Cooper K, Bendele MG, Cresawn SG. v6199 June 2025 https://observablehq.com/@cresawn-labs/cluster-report

Cresawn SG, Bogel M, Day N, Jacobs-Sera D, Hendrix RW, Hatfull GF. 2011. Phamerator: a bioinformatic tool for comparative bacteriophage genomics. BMC Bioinformatics 12: 10.1186/1471-2105-12-395. DOI: 10.1186/1471-2105-12-395

Delcher AL, Bratke KA, Powers EC, Salzberg SL. 2007. Identifying bacterial genes and endosymbiont DNA with Glimmer. Bioinformatics 23: 673-679. DOI: 10.1093/bioinformatics/btm009

Gordon D, Green P. 2013. *Consed:* a graphical editor for next-generation sequencing. Bioinformatics 29: 2936-2937. DOI: 10.1093/bioinformatics/btt515

Green SI, Clark JR, Santos HH, Weesner KE, Salazar KC, Aslam S, et al., Maresso. 2023. A Retrospective, Observational Study of 12 Cases of Expanded-Access Customized Phage Therapy: Production, Characteristics, and Clinical Outcomes. Clinical Infectious Diseases 77: 1079-1091. DOI: 10.1093/cid/ciad335

Hallgren J, Tsirigos KD, Pedersen MD, Almagro Armenteros JJ, Marcatili P, Nielsen H, Krogh A, Winther O. 2022. DeepTMHMM predicts alpha and beta transmembrane proteins using deep neural networks. : 10.1101/2022.04.08.487609. DOI: 10.1101/2022.04.08.487609

Hirokawa T, Boon-Chieng S, Mitaku S. 1998. SOSUI: classification and secondary structure prediction system for membrane proteins.. Bioinformatics 14: 378-379. DOI: 10.1093/bioinformatics/14.4.378

Jiang H, Lei R, Ding SW, Zhu S. 2014. Skewer: a fast and accurate adapter trimmer for next-generation sequencing paired-end reads. BMC Bioinformatics 15: 10.1186/1471-2105-15-182. DOI: 10.1186/1471-2105-15-182

Laslett D. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. Nucleic Acids Research 32: 11-16. DOI: 10.1093/nar/gkh152

Lowe TM, Eddy SR. 1997. tRNAscan-SE: A Program for Improved Detection of Transfer RNA Genes in Genomic Sequence. Nucleic Acids Research 25: 955-964. DOI: 10.1093/nar/25.5.0955

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal 17: 10. DOI: 10.14806/ej.17.1.200

Murphy C, Wilczak K, Frost VJ. 2025a. Unsuccessful Lysogen Formation for Subcluster C1 Phage, SVoro. HHMI Science Education Alliance (SEA) Faculty Group, QUBES Educational Resources. http://dx.doi.org/10.25334/JGK7-7Y18

Murphy C, Wilczak K, Frost VJ 2025b. Unsuccessful Lysogen Formation for Subcluster B2 Phage, PrinceCharming. HHMI Science Education Alliance (SEA) Faculty Group, QUBES Educational Resources. http://dx.doi.org/10.25334/RP14-2321

Chinemerem Nwobodo D, Ugwu MC, Oliseloke Anie C, Al-Ouqaili MTS, Chinedu Ikem J, Victor Chigozie U, Saki M. 2022. Antibiotic resistance: The challenges and some emerging strategies for tackling a global menace. Journal of Clinical Laboratory Analysis 36: 10.1002/jcla.24655. DOI: 10.1002/jcla.24655

Pope WH, Jacobs-Sera D. 2017. Annotation of Bacteriophage Genome Sequences Using DNA Master: An Overview. Methods in Molecular Biology,Bacteriophages : 217-229. DOI: 10.1007/978-1-4939-7343-9_16

Pope WH, Mavrich TN, Garlena RA, Guerrero-Bustamante CA, Jacobs-Sera D, Montgomery MT, et al., Hatfull. 2017. Bacteriophages of *Gordonia* spp. Display a Spectrum of Diversity and Genetic Relationships. mBio 8: 10.1128/mbio.01069-17. DOI: 10.1128/mBio.01069-17

Rinehart CA, Gaffney B, Wood JD, Smith S. 2016. PECAAN, a Phage Evidence Collection And Annotation Network. https://discover.kbrinsgd.org/login

Russell DA, Hatfull GF. 2016. PhagesDB: the actinobacteriophage database. Bioinformatics 33: 784-786. DOI: 10.1093/bioinformatics/btw711

Soding J, Biegert A, Lupas AN. 2005. The HHpred interactive server for protein homology detection and structure prediction. Nucleic Acids Research 33: W244-W248. DOI: 10.1093/nar/gki408

Teney Cc, Poupelin JC, Briot T, Le Bouar M, Fevre C, Brosset S, et al., Ferry. 2024. Phage Therapy in a Burn Patient Colonized with Extensively Drug-Resistant *Pseudomonas aeruginosa* Responsible for Relapsing Ventilator-Associated Pneumonia and Bacteriemia. Viruses 16: 1080. DOI: 10.3390/v16071080

Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. PLOS Computational Biology 13: e1005595. DOI: 10.1371/journal.pcbi.1005595

Yang Y, Tan X, Xiong M, Liu Z, Lu S, Ge H, et al., Pan. 2025. A case report of bacteriophage therapy for the treatment of lung infection due to carbapenem-resistant *Pseudomonas aeruginosa*. Scientific Reports 15: 10.1038/s41598-025-17510-3. DOI: 10.1038/s41598-025-17510-3

Zorawik M, Jacobs-Sera D, Freise AC, SEA-PHAGES, Reddi K. 2024. Isolation of Bacteriophages on Actinobacteria Hosts. Methods in Molecular Biology,Phage Engineering and Analysis : 273-298. DOI: 10.1007/978-1-0716-3798-2_17

Funding: V.F. acknowledges support by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number P20GM103499. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflicts of Interest: The authors declare that there are no conflicts of interest present.

Author Contributions: Kristi M Westover: supervision, investigation, validation, writing - original draft, writing - review editing. Kailyn C Cabana: investigation, writing - original draft, writing - review editing. Simon SE Cherry: investigation, writing - original draft, writing - review editing. Norah B Kay: investigation, writing - original draft, writing - review editing. Saachi Mehra: investigation, writing - original draft, writing - review editing. Olivia L Morgan: investigation, writing - original draft, writing - review editing. Ash H Olivet: investigation, writing - original draft, writing - review editing. Samuel W Reese: investigation, writing - original draft, writing - review editing. Sasha J Vorontsov: investigation, writing - original draft, writing - review editing. Ella A Williams: investigation, writing - original draft, writing - review editing. Mohammed RS Zyara: investigation, writing - original draft, writing - review editing. Victoria J Frost: supervision, investigation, validation, writing - original draft, writing - review editing.

Reviewed By: Anonymous

History: Received August 4, 2026 **Revision Received** October 26, 2026 **Accepted** November 3, 2026 **Published Online** November 10, 2026 **Indexed** November 24, 2026

Copyright: © 2026 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Westover KM, Cabana KC, Cherry SS, Kay NB, Mehra S, Morgan OL, et al., Frost VJ. 2026. Complete Genome Sequences of *Mycobacterium smegmatis* Phages SVoro and PrinceCharming . microPublication Biology. 10.17912/micropub.biology.002324